

Mapping Physical Tumor Microenvironment and Drug Delivery

Hadi T. Nia, Lance L. Munn, and Rakesh K. Jain



The physical microenvironment of pancreatic tumors is highly abnormal, and this causes significant challenges for drug delivery through multiple mechanisms. Measurements of tissue elasticity could be used as a physical

biomarker to assess aberrant drug delivery, and potentially guide stroma-targeting treatment strategies and patient stratification.

See related article by Wang et al.

In this issue of *Clinical Cancer Research*, Wang and colleagues demonstrate that shear modulus, a measure of tissue stiffness measured using elastography, correlates with drug delivery in two orthotopic mouse models of pancreatic tumors (1). Although increased matrix density can cause increased stiffness and hinder drug delivery, there are other aspects of the physical tumor microenvironment (PhysTME) that also affect drug delivery in desmoplastic tumors. To give the reader a more accurate view of the complex interplay between the PhysTME and drug delivery, we will start with a brief overview of the key elements of abnormal PhysTME in desmoplastic tumors, describe their etiology, and discuss the mechanisms underlying aberrant drug delivery in tumors. We will also discuss the translational and clinical relevance of elastography, one of the very few noninvasive modalities for characterizing the PhysTME, and discuss the potential use of elastography as a surrogate for abnormal physical properties in tumors.

The three main physical abnormalities in most desmoplastic tumors, including pancreatic adenocarcinomas, are as follows (2): (i) increased stiffness (also known as rigidity and elasticity), (ii) elevated solid stresses, and (iii) elevated interstitial fluid pressure (IFP). Stiffness or rigidity, defined as the resistance to deformation, is the first tumor mechanical abnormality to be recognized, and is the basis for well-established diagnostic and prognostic methods for a variety of cancer types. The biological processes that contribute to increased stiffness are deposition of extracellular matrix (ECM) and increased cross-linking of ECM constituents. Relevant to this discussion, matrix deposition and crosslinking tend to reduce the amount of space in the tissue available for movement of drugs and other materials (i.e., "pore size" ref. 3), thus restricting drug access into the tumor. It is well known that dense tissues are stiffer and have lower drug diffusion rates (Fig. 1; ref. 3); Wang and colleagues have, for the first time, confirmed this relationship by mapping drug delivery and tissue elasticity.

Tissues can be stiff or soft, but they can also have regions that are under tension or compression. Tension and compression are mechanical solid stresses, the result of forces contained and

transmitted by solid components of the tissue including the ECM and cellular elements. Solid stresses are elevated in many fibrotic tumors, but are nearly zero in the corresponding normal tissues (2, 4). Solid stresses are caused by cell proliferation or infiltration, matrix deposition, gel swelling, and resistance to the volume expansion from within and outside the tumor through the ECM and cell adhesion. Solid stresses impede drug delivery by compressing blood and lymphatic vessels (Fig. 1; ref. 2). When solid stress is reduced via antifibrotic treatments, drug delivery increases through decompression of blood vessels (5). This results in an improvement in tumor response and overall survival when antifibrotic strategies are combined with anticancer treatments [refs. 2, 5; Murphy and colleagues: A phase II study of neoadjuvant FOLFIRINOX in combination with losartan followed by chemoradiotherapy in locally advanced pancreatic cancer: R0 resection rate and clinical outcomes (submitted)].

Elevated stiffness and pent-up solid stresses are structural/mechanical properties of the tumor; in addition, there are abnormalities in the fluid environment of tumors that interfere with drug delivery. Normally, fluid travels easily through tissue between the blood and lymphatic systems, maintaining a fluid pressure close to zero in the interstitium. However, there are no functional lymphatic vessels in tumors, and the blood vessels are abnormally leaky. Thus, IFP is elevated throughout the tumor, only decreasing at the tumor periphery where functional lymphatics exist. The elevated IFP is approximately uniform across the tumor volume, with an abrupt drop at the tumor-normal tissue interface. The high uniform IFP across the tumor impedes the transport of drug and nutrients from the blood vessels within tumors. The sharp drop in IFP at the tumor edge also causes fluid to "ooze" into the peritumor normal tissue that promotes "washing" of the drug along with growth factors and cytokines from the tumor into the normal surrounding tissue (2).


Because of the well-established links between drug delivery and the PhysTME (Fig. 1), efforts to characterize tumor mechanopathologies have intensified. Although there are several approaches to quantify the three major components of the PhysTME, the only one that can currently be measured noninvasively is stiffness. Wang and colleagues proposed using shear wave elastography to map the shear modulus of pancreatic tumors and examine how well the shear modulus correlates with drug delivery in pancreatic tumors. This work is one of the first studies to provide a noninvasive measurement for a physical biomarker to correlate with drug delivery. Wang and colleagues created orthotopic human xenograft models of pancreatic cancer by implanting human cells in

Edwin L. Steele Laboratories, Department of Radiation Oncology, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts.

Corresponding Author: Rakesh K. Jain, Massachusetts General Hospital, Harvard Medical School, 100 Blossom Street, Cox 7, Boston, MA 02114. Phone: 617-726-4083; Fax: 617-724-1819; E-mail: jain@steele.mgh.harvard.edu

doi: 10.1158/1078-0432.CCR-18-3724

©2019 American Association for Cancer Research

**Figure 1.**

Drug delivery in tumors is affected by the physical microenvironment. Elevated fluid pressure, solid stress, and stiffness, the three main features of the physical microenvironment in most tumors with distinct origins and consequences, affect drug delivery through different mechanisms.

immune-deficient mice. Although a syngeneic model would have recapitulated the highly desmoplastic human pancreatic tumors more closely, the authors verified the presence of the key ECM components: collagen (via trichome staining) and hyaluronic acid (HA). The delivery of verteporfin, an FDA-approved fluorescent drug, was mapped in the tumor after intravenous perfusion. An important aspect of the study is the correlation of the spatial maps of drug delivery and ECM content with those of shear modulus. They perform these correlations within each tumor as well as globally, that is, across mice. A key finding is that shear modulus inversely correlates with drug uptake, when compared globally. As

expected, shear modulus was shown to correlate strongly with collagen both globally and locally. Interestingly, the correlation between shear modulus and HA has been weak when compared globally; however, they seem to correlate reasonably well when compared locally.

The main question that Wang and coworkers attempted to answer was whether stiffness is an appropriate biomarker to predict drug delivery. Another question indirectly addressed in this study was whether stiffness measurements are related to solid stress levels. Although increased matrix deposition is a common source of stiffness and solid stress, solid stress is also accumulated through other mechanisms and is dramatically affected by growth

pattern. For example, two tumors generated from the same pancreatic cancer cells might have similar stiffness values, but the solid stress levels can be significantly different (4).

In light of the discussion above, drug delivery is affected by three distinct abnormalities of the PhysTME, all based on different biological and/or physical mechanisms, including transport within blood vessels, transport from vessels into tissue, and transport through the interstitium. Matrix densification, which is reflected in increased stiffness, affects the interstitial transport outside the blood vessels. Although important, this does not reveal the whole story. To rigorously predict drug delivery, we would also need to know whether solid stresses are restricting local fluid flow by compressing blood or lymphatic vessels. In addition, we would need to know whether the drug can easily move across the vessel walls and whether IFP is elevated. Therefore, while tumor tissue elastography indicates how dense the matrix is, it cannot assess other properties that affect drug delivery in tumors. For a more accurate estimation of drug delivery, measurement of both solid stress and IFP would be needed.

Fortunately, elastography may be further extended to help fill this unmet need. Conventional elastography, which is based on linear elasticity, estimates stiffness, which does not necessarily predict solid stress in tumors. However, by utilizing the strain-stiffening mechanism, a nonlinear effect that describes how stiff a tissue becomes under tension or compression, it may be possible to provide a measure of solid stress using modifications of elastography. Ongoing research into the strain-stiffening phenomenon may result in new elastography modalities to decouple stiffness from solid stress and provide a noninvasive measure of solid stress in tumors.

Another challenge is to provide elasticity maps of pancreatic tumors *in situ*. Wang and colleagues performed the measurement *ex vivo* with tumors embedded in a gel for a higher quality of

imaging, as extending the methodology to measuring the shear modulus deep in patient tissues is not straightforward. Tumors such as breast and brain are relatively accessible, and thus, shear wave elastography can be performed *in vivo*. However, pancreatic tumors are deeper in the abdomen, and the surrounding organs confound the use of shear wave elastography. Currently, elastography in pancreatic tumors is limited to endoscopic ultrasound elastography, which provides a relative map of stiffness within a tumor, and therefore cannot be used longitudinally or to compare results among patients. Noninvasive, large-scale quantification of tumor stiffness, which can be used to compare stiffness of tumors among patients, is an unmet need with high clinical potential for improved detection, diagnosis, and design of antifibrotic treatment strategies for pancreatic and other desmoplastic cancers.

Disclosure of Potential Conflicts of Interest

R.K. Jain serves on the board of trustees for Tekla Healthcare Investors, Tekla Life Sciences Investors, Tekla Healthcare Opportunities Fund, and Tekla World Healthcare Fund, reports receiving speakers bureau honoraria from Amgen, holds ownership interest in Enlight, Ophthotech, and SynDevRx, and is a consultant/advisory board member for Merck, Ophthotech, Pfizer, SPARC, SynDevRx, and XTuit. No potential conflicts of interest were disclosed by the other authors.

Acknowledgments

This work was supported by the NCI (grant numbers P01-CA080124, P50-CA165962, R01-CA129371, R01-CA208205, and U01-CA 224348; to R.K. Jain), the NCI Outstanding Investigator Award (grant number R35-CA197743; to R.K. Jain), the Lustgarten Foundation, the Ludwig Center at Harvard, the National Foundation for Cancer Research, and the Gates Foundation (to R.K. Jain), R01-HL128168 (to L.L. Munn), and F32-CA216944-01 (to H.T. Nia).

Received December 3, 2018; revised December 30, 2018; accepted January 7, 2019; published first January 10, 2019.

References

1. Wang H, Mislati R, Ahmed R, Vincent P, Nwabunwanne SF, Gunn JR, et al. Elastography can map the local inverse relationship between shear modulus and drug delivery within the pancreatic ductal adenocarcinoma microenvironment. *Clin Cancer Res* 2018 Oct 23 [Epub ahead of print].
2. Jain RK, Martin JD, Stylianopoulos T. The role of mechanical forces in tumor growth and therapy. *Annu Rev Biomed Eng* 2014; 16:321.
3. Netti PA, Berk DA, Swartz MA, Grodzinsky AJ, Jain RK. Role of extracellular matrix assembly in interstitial transport in solid tumors. *Cancer Res* 2000; 60:2497–503.
4. Nia HT, Liu H, Seano G, Datta M, Jones D, Rahbari N, et al. Solid stress and elastic energy as measures of tumour mechanopathology. *Nat Biomed Eng* 2016;1:0004.
5. Chauhan VP, Martin JD, Liu H, Lacorre DA, Jain SR, Kozin SV, et al. Angiotensin inhibition enhances drug delivery and potentiates chemotherapy by decompressing tumour blood vessels. *Nat Commun* 2013;4:2516.

Clinical Cancer Research

Mapping Physical Tumor Microenvironment and Drug Delivery

Hadi T. Nia, Lance L. Munn and Rakesh K. Jain

Clin Cancer Res Published OnlineFirst January 10, 2019.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-18-3724](https://doi.org/10.1158/1078-0432.CCR-18-3724)

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link <http://clincancerres.aacrjournals.org/content/early/2019/02/05/1078-0432.CCR-18-3724>. Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.